

Analysis & Numerical Treatment of Singular IVPs

Part II

O. Koch, P. Kofler and Ewa B. Weinmüller,
Institute for Applied Mathematics and Numerical Analysis,
University of Technology, Vienna, Austria.

Contents

- Singular IVPs
- Implicit Euler method
- Asymptotic error expansion for the approximation computed by the implicit Euler method
- Acceleration technique
 - Zadunaisky's idea
 - Iterated Defect Correction (IDeC)
 - Experimental results
- Conclusion

Singular IVPs

$$z'(t) = \frac{M(t)}{t}z(t) + f(t, z(t)), t \in (0, 1]$$

- $M(t) = M + tC(t)$, $C \in C[0, 1]$
- $\lambda(M) : \operatorname{Re}(\lambda) < 0$ or $\lambda = 0$
- $f(t, y) \in C([0, 1] \times \mathbb{R}^n)$

$$Mz(0) = 0 \quad (\iff z \in C[0, 1])$$

- yields $\operatorname{rank}(M) = n - m$ equations

$$B_0z(0) = \beta$$

- $B_0 \in \mathbb{R}^{(m \times n)}$, $\beta \in \mathbb{R}^m$

Singular IVPs – theoretical results

- $C \in C^p[0, 1], p \geq 0$

- $f(t, y) \in C^p([0, 1] \times \mathbb{R}^n),$

Lipschitz-condition with respect to y

- $m \times m$ -matrix $B_0 R$ nonsingular,

R projection onto eigenspace of M


for eigenvalue $\lambda = 0$



existence and uniqueness of solution z

$$z \in C^{p+1}[0, 1] \quad (p = 6)$$

Implicit Euler method



$$\Delta_h = (t_0, t_1, \dots, t_i, \dots, t_N), \quad N = 1/h$$

$$\left\{ \begin{array}{l} \frac{v_i - v_{i-1}}{h} = \frac{M(t_i)}{t_i} v_i + f(t_i, v_i), \quad i = 1(1)N \\ B_0 v_0 = \beta \\ M v_0 = 0 \end{array} \right.$$

$$v_h = (v_0, v_1, \dots, v_i, \dots, v_N)$$

⇓

$$|v_i - z(t_i)| = O(h), \quad i = 0(1)N, \quad \text{for } h \rightarrow 0$$

Asymtotic error expansion

- $v_h = (v_0, \dots, v_i, \dots, v_N)$ grid vector
solution obtained by implicit Euler
method on mesh Δ_h

- we make an ansatz

$$v_i = z(t_i) + \sum_{j=1}^5 h^j e_j(t_i) + r_i, t_i \in \Delta_h$$

- and have to show

– $\exists e_j$ smooth

– $|r_i| = O(h^6)$, $i = 0(1)N$, for $h \rightarrow 0$

Variational equations

singular IVPs

$$\begin{cases} e_1'(t) - \frac{M(t)}{t}e_1(t) = f_y(t, z(t))e_1(t) + \frac{1}{2}z''(t) \\ e_1(0) = 0 \\ e_2'(t) - \frac{M(t)}{t}e_2(t) = f_y(t, z(t))e_2(t) + \frac{1}{2}e_1''(t) + \\ \quad \frac{1}{2}f_{yy}(t, z(t))e_1^2(t) - \frac{1}{6}z^{(3)}(t) \\ e_2(0) = 0 \end{cases}$$

$$z \in C^7[0, 1]$$

\Downarrow

$$e_1 \in C^6[0, 1] \implies e_2 \in C^5[0, 1]$$

\Downarrow

e_j smooth for all further j , $3 \leq j \leq 5$

Remainder term

- r_i , $i = 1(1)N$, satisfies

$$\begin{cases} \frac{r_i - r_{i-1}}{h} = \frac{M(t_i)}{t_i} r_i + g(t_i, r_i) + l_i \\ r_0 = 0 \end{cases}$$

- with

$$g(t_i, r_i) := \int_0^1 f_y(t_i, v_i - (1 - \tau)r_i) d\tau \cdot r_i$$

and $l_i = O(h^6)$

- by a contraction argument we show that there exists an $L < 1$ with

$$|r_i| \leq \text{const} \frac{1}{1 - L} |l_i| = O(h^6), \text{ for } h \rightarrow 0$$

Zadunaisky's idea

- solve

$$\begin{cases} z'(t) = F(t, z(t)), t \in (0, 1] \\ z(0) = \beta \end{cases}$$

by implicit Euler method

- obtain grid vector solution

$$z_h^{(0)} = (z_0^{(0)}, \dots, z_i^{(0)}, \dots, z_N^{(0)})$$

- interpolate $(t_0, z_0^{(0)}), \dots, (t_N, z_N^{(0)})$ by polynomial $p^{(0)}(t)$ of degree $m = N$

- solve “neighboring problem”

$$\begin{cases} z'(t) = F(t, z(t)) + p^{(0)'}(t) - F(t, p^{(0)}(t)) \\ z(0) = p^{(0)}(0) = \beta \end{cases}$$

Zadunaisky's idea – continued

- obtain grid vector solution $p_h^{(0)}$ by implicit Euler method on the same grid Δ_h
- use known global error $z_h^{(0)} - p_h^{(0)}$ as estimation for the unknown global error of the original problem
- accuracy by the heuristic argumentation
- implementation:
 - continuous function $p^{(0)}(t)$ composed of polynomials with degree $m < N$

Iterated Defect Correction (IDeC)

- improve approximation

$$z_h^{(1)} := z_h^{(0)} + (z_h^{(0)} - p_h^{(0)})$$

- interpolate $(t_0, z_0^{(1)}), \dots, (t_N, z_N^{(1)})$ by new polynomial $p^{(1)}(t)$

- solve “neighboring problem” with defect

$$p^{(1)'}(t) - F(t, p^{(1)}(t))$$

- obtain grid vector solution $p_h^{(1)}$ by implicit Euler method on the same grid Δ_h

- improve approximation again

$$z_h^{(2)} := z_h^{(0)} + (z_h^{(1)} - p_h^{(1)})$$

IDeC – expectations

✓ practical basis for IDeC:

Zadunaisky's and Stetter's idea

$$n := \min(m, 6)$$

$$|z_{i_h^*}^{(j-1)} - z(t_*)| = O(h^j), \text{ for } 1 \leq j \leq n$$

$$|z_{i_h^*}^{(j-1)} - z(t_*)| = O(h^n), \text{ for } j > n$$

✓ theoretical basis for IDeC:

asymptotic error expansion

“Emden differential equation”

$$\begin{cases} z'(t) = \frac{1}{t} \begin{pmatrix} 0 & 1 \\ 0 & -1 \end{pmatrix} z(t) - t \begin{pmatrix} 0 \\ z_1^5(t) \end{pmatrix} \\ z(0) = (1, 0)^T \end{cases}$$

| h | $\ \varepsilon_h\ _h$ | p | c | $\ \varepsilon_h^{(1)}\ _h$ | $p^{(1)}$ | $c^{(1)}$ |
|---------------------|-----------------------|-------|-----------------------|-----------------------------|-----------|-----------------------|
| $1/5$ | $2.4 \cdot 10^{-02}$ | 0.834 | $-9.2 \cdot 10^{-01}$ | $7.7 \cdot 10^{-02}$ | 1.835 | $-1.4 \cdot 10^{-01}$ |
| $1/5 \cdot 2^{-1}$ | $1.3 \cdot 10^{-02}$ | 0.921 | $-1.1 \cdot 10^{-01}$ | $2.1 \cdot 10^{-03}$ | 1.918 | $-1.7 \cdot 10^{-01}$ |
| $1/5 \cdot 2^{-2}$ | $7.1 \cdot 10^{-02}$ | 0.960 | $-1.2 \cdot 10^{-01}$ | $5.7 \cdot 10^{-03}$ | 1.959 | $-2.0 \cdot 10^{-01}$ |
| $1/5 \cdot 2^{-3}$ | $3.6 \cdot 10^{-02}$ | 0.980 | $-1.3 \cdot 10^{-01}$ | $1.4 \cdot 10^{-04}$ | 1.979 | $-2.1 \cdot 10^{-01}$ |
| $1/5 \cdot 2^{-4}$ | $1.8 \cdot 10^{-03}$ | 0.990 | $-1.4 \cdot 10^{-01}$ | $3.7 \cdot 10^{-04}$ | 1.989 | $-2.2 \cdot 10^{-01}$ |
| $1/5 \cdot 2^{-5}$ | $9.3 \cdot 10^{-03}$ | 0.995 | $-1.4 \cdot 10^{-01}$ | $9.4 \cdot 10^{-05}$ | 1.994 | $-2.3 \cdot 10^{-01}$ |
| $1/5 \cdot 2^{-6}$ | $4.7 \cdot 10^{-03}$ | 0.997 | $-1.4 \cdot 10^{-01}$ | $2.3 \cdot 10^{-06}$ | 1.997 | $-2.3 \cdot 10^{-01}$ |
| $1/5 \cdot 2^{-7}$ | $2.3 \cdot 10^{-04}$ | 0.998 | $-1.4 \cdot 10^{-01}$ | $5.9 \cdot 10^{-06}$ | 1.998 | $-2.4 \cdot 10^{-01}$ |
| $1/5 \cdot 2^{-8}$ | $1.1 \cdot 10^{-04}$ | 0.999 | $-1.5 \cdot 10^{-01}$ | $1.4 \cdot 10^{-07}$ | 1.999 | $-2.4 \cdot 10^{-01}$ |
| $1/5 \cdot 2^{-9}$ | $5.8 \cdot 10^{-04}$ | 0.999 | $-1.5 \cdot 10^{-01}$ | $3.7 \cdot 10^{-07}$ | 1.999 | $-2.4 \cdot 10^{-01}$ |
| $1/5 \cdot 2^{-10}$ | $2.9 \cdot 10^{-05}$ | 0.999 | $-1.5 \cdot 10^{-01}$ | $9.2 \cdot 10^{-08}$ | 2.000 | $-2.4 \cdot 10^{-01}$ |

| h | $\ \varepsilon_h^{(2)}\ _h$ | $p^{(2)}$ | $c^{(2)}$ | $\ \varepsilon_h^{(3)}\ _h$ | $p^{(3)}$ | $c^{(3)}$ |
|--------------------|-----------------------------|-----------|-----------------------|-----------------------------|-----------|-----------------------|
| 1/5 | $1.6 \cdot 10^{-03}$ | 2.924 | $-1.8 \cdot 10^{-01}$ | $9.5 \cdot 10^{-03}$ | 3.330 | $-2.0 \cdot 10^{-01}$ |
| $1/5 \cdot 2^{-1}$ | $2.1 \cdot 10^{-04}$ | 2.918 | $-1.7 \cdot 10^{-01}$ | $9.5 \cdot 10^{-04}$ | 3.525 | $-3.1 \cdot 10^{+00}$ |
| $1/5 \cdot 2^{-2}$ | $2.8 \cdot 10^{-05}$ | 2.731 | $-1.0 \cdot 10^{-01}$ | $8.2 \cdot 10^{-05}$ | 3.770 | $-6.6 \cdot 10^{+00}$ |
| $1/5 \cdot 2^{-3}$ | $4.3 \cdot 10^{-05}$ | 2.868 | $-1.6 \cdot 10^{-01}$ | $6.0 \cdot 10^{-06}$ | 3.889 | $-1.0 \cdot 10^{+00}$ |
| $1/5 \cdot 2^{-4}$ | $5.9 \cdot 10^{-06}$ | 2.934 | $-2.2 \cdot 10^{-01}$ | $4.0 \cdot 10^{-07}$ | 3.946 | $-1.3 \cdot 10^{+00}$ |
| $1/5 \cdot 2^{-5}$ | $7.7 \cdot 10^{-07}$ | 2.967 | $-2.6 \cdot 10^{-01}$ | $2.6 \cdot 10^{-09}$ | 3.973 | $-1.5 \cdot 10^{+00}$ |
| $1/5 \cdot 2^{-6}$ | $9.8 \cdot 10^{-08}$ | 2.983 | $-2.9 \cdot 10^{-01}$ | $1.6 \cdot 10^{-10}$ | 3.974 | $-1.5 \cdot 10^{+00}$ |
| $1/5 \cdot 2^{-7}$ | $1.2 \cdot 10^{-09}$ | 2.990 | $-3.0 \cdot 10^{-01}$ | $1.0 \cdot 10^{-11}$ | 3.613 | $-1.4 \cdot 10^{-01}$ |
| $1/5 \cdot 2^{-8}$ | $1.5 \cdot 10^{-10}$ | 2.979 | $-2.8 \cdot 10^{-01}$ | $8.7 \cdot 10^{-12}$ | 0.958 | $-8.3 \cdot 10^{-09}$ |
| h | $\ \varepsilon_h^{(4)}\ _h$ | $p^{(4)}$ | $c^{(4)}$ | $\ \varepsilon_h^{(5)}\ _h$ | $p^{(5)}$ | $c^{(5)}$ |
| 1/5 | $8.1 \cdot 10^{-03}$ | 4.954 | $-2.3 \cdot 10^{+00}$ | $5.4 \cdot 10^{-03}$ | 5.558 | $-4.1 \cdot 10^{+01}$ |
| $1/5 \cdot 2^{-1}$ | $2.6 \cdot 10^{-05}$ | 5.079 | $-3.1 \cdot 10^{+00}$ | $1.1 \cdot 10^{-05}$ | 4.921 | $-9.5 \cdot 10^{+00}$ |
| $1/5 \cdot 2^{-2}$ | $7.7 \cdot 10^{-06}$ | 5.095 | $-3.3 \cdot 10^{+01}$ | $3.7 \cdot 10^{-06}$ | 5.141 | $-1.8 \cdot 10^{+00}$ |
| $1/5 \cdot 2^{-3}$ | $2.2 \cdot 10^{-08}$ | 5.084 | $-3.1 \cdot 10^{+01}$ | $1.0 \cdot 10^{-08}$ | 5.102 | $-1.6 \cdot 10^{+00}$ |
| $1/5 \cdot 2^{-4}$ | $6.6 \cdot 10^{-09}$ | 4.995 | $-2.1 \cdot 10^{+00}$ | $3.1 \cdot 10^{-10}$ | 5.080 | $-1.4 \cdot 10^{+00}$ |
| $1/5 \cdot 2^{-5}$ | $2.0 \cdot 10^{-11}$ | 4.881 | $-1.2 \cdot 10^{+00}$ | $9.2 \cdot 10^{-11}$ | 5.395 | $-7.2 \cdot 10^{+01}$ |
| $1/5 \cdot 2^{-6}$ | $7.1 \cdot 10^{-12}$ | 2.400 | $-7.3 \cdot 10^{-06}$ | $2.1 \cdot 10^{-13}$ | 0.770 | $-1.8 \cdot 10^{-11}$ |

Example 2

$$\begin{cases} z'(t) = \frac{1}{t} \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} z(t) - t \begin{pmatrix} 0 \\ e^{z_1(t)} - \frac{B^2 - 6B + 1}{Bt^2 + 1} \end{pmatrix} \\ z(0) = (2 \ln(B + 1), 0)^T, \quad B = 3 + \sqrt{8} \end{cases}$$

| h | $\ \varepsilon_h\ _h$ | p | c | $\ \varepsilon_h^{(1)}\ _h$ | $p^{(1)}$ | $c^{(1)}$ |
|---------------------|-----------------------|-------|-----------------------|-----------------------------|-----------|-----------------------|
| $1/5 \cdot 2^{-1}$ | 4.9 | 0.608 | $-2.0 \cdot 10^{+00}$ | $3.1 \cdot 10^{-01}$ | 1.385 | $-7.5 \cdot 10^{+01}$ |
| $1/5 \cdot 2^{-2}$ | 3.2 | 0.783 | $-3.4 \cdot 10^{+01}$ | $1.1 \cdot 10^{-01}$ | 1.682 | $-1.8 \cdot 10^{+01}$ |
| $1/5 \cdot 2^{-3}$ | $1.8 \cdot 10^{-01}$ | 0.886 | $-4.9 \cdot 10^{+01}$ | $3.7 \cdot 10^{-01}$ | 1.838 | $-3.2 \cdot 10^{+02}$ |
| $1/5 \cdot 2^{-4}$ | $1.0 \cdot 10^{-01}$ | 0.941 | $-6.3 \cdot 10^{+01}$ | $1.0 \cdot 10^{-02}$ | 1.917 | $-4.6 \cdot 10^{+02}$ |
| $1/5 \cdot 2^{-5}$ | $5.3 \cdot 10^{-01}$ | 0.970 | $-7.3 \cdot 10^{+01}$ | $2.7 \cdot 10^{-03}$ | 1.958 | $-5.6 \cdot 10^{+02}$ |
| $1/5 \cdot 2^{-6}$ | $2.7 \cdot 10^{-02}$ | 0.985 | $-8.0 \cdot 10^{+01}$ | $7.0 \cdot 10^{-03}$ | 1.979 | $-6.4 \cdot 10^{+02}$ |
| $1/5 \cdot 2^{-7}$ | $1.3 \cdot 10^{-02}$ | 0.992 | $-8.3 \cdot 10^{+01}$ | $1.7 \cdot 10^{-04}$ | 1.989 | $-6.8 \cdot 10^{+02}$ |
| $1/5 \cdot 2^{-8}$ | $6.9 \cdot 10^{-02}$ | 0.996 | $-8.6 \cdot 10^{+01}$ | $4.5 \cdot 10^{-04}$ | 1.994 | $-7.1 \cdot 10^{+02}$ |
| $1/5 \cdot 2^{-9}$ | $3.4 \cdot 10^{-02}$ | 0.998 | $-8.7 \cdot 10^{+01}$ | $1.1 \cdot 10^{-05}$ | 1.997 | $-7.2 \cdot 10^{+02}$ |
| $1/5 \cdot 2^{-10}$ | $1.7 \cdot 10^{-03}$ | 0.999 | $-8.8 \cdot 10^{+01}$ | $2.8 \cdot 10^{-06}$ | 1.998 | $-7.3 \cdot 10^{+02}$ |

| h | $\ \varepsilon_h^{(2)}\ _h$ | $p^{(2)}$ | $c^{(2)}$ | $\ \varepsilon_h^{(3)}\ _h$ | $p^{(3)}$ | $c^{(3)}$ |
|--------------------|-----------------------------|-----------|-----------------------|-----------------------------|-----------|-----------------------|
| $1/5 \cdot 2^{-1}$ | $1.7 \cdot 10^{-01}$ | 2.481 | $-5.1 \cdot 10^{+02}$ | $7.1 \cdot 10^{-01}$ | 3.125 | $-9.5 \cdot 10^{+02}$ |
| $1/5 \cdot 2^{-2}$ | $3.0 \cdot 10^{-02}$ | 2.895 | $-1.7 \cdot 10^{+02}$ | $8.2 \cdot 10^{-02}$ | 3.107 | $-9.0 \cdot 10^{+02}$ |
| $1/5 \cdot 2^{-3}$ | $4.1 \cdot 10^{-02}$ | 3.003 | $-2.6 \cdot 10^{+02}$ | $9.5 \cdot 10^{-03}$ | 3.524 | $-4.2 \cdot 10^{+03}$ |
| $1/5 \cdot 2^{-4}$ | $5.1 \cdot 10^{-03}$ | 2.977 | $-2.3 \cdot 10^{+02}$ | $8.2 \cdot 10^{-04}$ | 3.806 | $-1.4 \cdot 10^{+03}$ |
| $1/5 \cdot 2^{-5}$ | $6.5 \cdot 10^{-04}$ | 2.988 | $-2.5 \cdot 10^{+02}$ | $5.9 \cdot 10^{-05}$ | 3.915 | $-2.5 \cdot 10^{+03}$ |
| $1/5 \cdot 2^{-6}$ | $8.2 \cdot 10^{-05}$ | 2.993 | $-2.6 \cdot 10^{+02}$ | $3.9 \cdot 10^{-06}$ | 3.960 | $-3.2 \cdot 10^{+04}$ |
| $1/5 \cdot 2^{-7}$ | $1.0 \cdot 10^{-06}$ | 2.996 | $-2.6 \cdot 10^{+02}$ | $2.5 \cdot 10^{-08}$ | 3.980 | $-3.7 \cdot 10^{+04}$ |
| $1/5 \cdot 2^{-8}$ | $1.2 \cdot 10^{-07}$ | 2.998 | $-2.6 \cdot 10^{+02}$ | $1.5 \cdot 10^{-09}$ | 3.979 | $-3.7 \cdot 10^{+04}$ |
| $1/5 \cdot 2^{-9}$ | $1.6 \cdot 10^{-08}$ | 3.001 | $-2.7 \cdot 10^{+02}$ | $1.0 \cdot 10^{-10}$ | 3.666 | $-3.1 \cdot 10^{+03}$ |
| h | $\ \varepsilon_h^{(4)}\ _h$ | $p^{(4)}$ | $c^{(4)}$ | $\ \varepsilon_h^{(5)}\ _h$ | $p^{(5)}$ | $c^{(5)}$ |
| $1/5 \cdot 2^{-1}$ | $3.0 \cdot 10^{-02}$ | 1.719 | $-1.5 \cdot 10^{+00}$ | $5.3 \cdot 10^{-01}$ | 2.829 | $-3.6 \cdot 10^{+02}$ |
| $1/5 \cdot 2^{-2}$ | $9.2 \cdot 10^{-02}$ | 4.818 | $-1.7 \cdot 10^{+04}$ | $7.5 \cdot 10^{-02}$ | 6.257 | $-1.0 \cdot 10^{+06}$ |
| $1/5 \cdot 2^{-3}$ | $3.2 \cdot 10^{-03}$ | 5.328 | $-1.1 \cdot 10^{+05}$ | $9.8 \cdot 10^{-04}$ | 4.749 | $-3.9 \cdot 10^{+04}$ |
| $1/5 \cdot 2^{-4}$ | $8.1 \cdot 10^{-05}$ | 5.224 | $-7.1 \cdot 10^{+05}$ | $3.6 \cdot 10^{-05}$ | 5.272 | $-3.9 \cdot 10^{+05}$ |
| $1/5 \cdot 2^{-5}$ | $2.1 \cdot 10^{-07}$ | 4.955 | $-1.8 \cdot 10^{+04}$ | $9.4 \cdot 10^{-07}$ | 5.315 | $-4.9 \cdot 10^{+05}$ |
| $1/5 \cdot 2^{-6}$ | $7.0 \cdot 10^{-08}$ | 4.920 | $-1.4 \cdot 10^{+04}$ | $2.3 \cdot 10^{-09}$ | 5.256 | $-3.4 \cdot 10^{+05}$ |
| $1/5 \cdot 2^{-7}$ | $2.3 \cdot 10^{-10}$ | 4.985 | $-2.2 \cdot 10^{+04}$ | $6.2 \cdot 10^{-10}$ | 5.552 | $-2.3 \cdot 10^{+05}$ |

Conclusion

- order sequence is $O(h), O(h^2), O(h^3), \dots$
as expected
- IDeC is very efficient



further investigations:

- Is implicit middle-point rule $O(h^2)$?
- Exists an asymptotic error expansion?
- Has the IDeC the order sequence $O(h^2), O(h^4), O(h^6), \dots$?